

AUSTRALIAN MUSEUM SCIENTIFIC PUBLICATIONS

Ogilby, J. D., 1893. Description of a new shark from the Tasmanian coast.
Records of the Australian Museum 2(5): 62–63. [30 September 1893].

doi:10.3853/j.0067-1975.2.1893.1194

ISSN 0067-1975

Published by the Australian Museum, Sydney

nature culture **discover**

Australian Museum science is freely accessible online at
<http://publications.australianmuseum.net.au>
6 College Street, Sydney NSW 2010, Australia



common in New South Wales, *T. proximus* is, on the other hand, comparatively rare. Owing, however, to the very limited number of observations made upon the Australian *Typhlopidae*, it would at present be extremely unwise to hazard many remarks upon their distribution.

DESCRIPTION OF A NEW SHARK FROM THE TASMANIAN
COAST.

BY J. DOUGLAS OGILBY.

CENTRINA BRUNIENSIS, *sp. nov.*

Centrina bruniensis, Morton (*in lit.*)

Body oblong, with the back and sides rounded, and the belly flattened. Head small and strongly depressed, its breadth equal to the distance between the tip of the snout and the spiracle: snout short and obtuse, the distance between its tip and the nearest point of the mouth less than that between the same and the anterior margin of the eye. Nostrils equidistant from the eye and the extremity of the snout. Eye large, with a strong bony supraorbital ridge, situated midway between the tip of the snout and the anterior gill-opening. Spiracles large, opening behind the upper half of the eye, with a moderate intervening space. Mouth small and transverse, with the lateral groove very broad and deep. Upper jaw with a patch of small, conical, curved teeth anteriorly, consisting of about four irregular rows; a single series of much larger, erect, compressed, minutely serrated, scalpriform teeth in the lower jaw. Gill-openings small, the posterior one pierced immediately in front of the base of the pectoral fin. The first dorsal commences above the middle gill-opening, and rises by a continuous and equal gradation to the spine, its outer margin being straight; behind the spine the rise is much more abrupt, and the contour is slightly convex with the tip rounded; the posterior margin is deeply concave; the height of the fin beneath its extremity is equal to the distance between the anterior gill-opening and the tip of the snout, that of the spine equal to the head in front of the spiracle; the spine is situated in the anterior portion of the last fourth of the base of the fin, is perfectly

straight, with a slight inclination forwards, and protrudes a short distance beyond the membrane; its base is exactly midway between the tip of the snout and the origin of the caudal, while the distance between the bases of the two dorsal spines is but little more than the length of the base of the first dorsal in front of its spine, and five-sevenths of the length of the fish in front of it; the intradorsal ridge is very strongly developed; the second dorsal has a general resemblance in shape to the first, but is not so large; the upper margin is more regularly even, and the extremity, which is much more pointed, hangs vertically above the base of the caudal, instead of falling within the vertical from its own base, as with the anterior fin; the length of its base is equal to that of the intradorsal space, and to the height of the fin beneath its tip, and is four-sevenths of the outer margin; the spine is situated in the latter portion of the anterior half of the fin, and is gently curved backwards throughout its entire length; in height it is but little less than that of the first dorsal; the pectoral fin is well developed and pointed, its length equal to the space which divides its anterior basal margin from the nostril; the distance between its base and that of the ventral is two-fifths longer than that between the dorsal spines, and is traversed by a strongly developed lateral ridge; the ventral fin commences beneath the spine of the second dorsal, and the distance between its termination and the origin of the lower caudal lobe is equal to that between the second dorsal and the caudal fin; the caudal lobes are well developed; the outer margin of the upper lobe is straight, the angle and the posterior margin rounded; the lower lobe is triangular, with the anterior margin slightly concave, and equal in length to the posterior margin, which is sinuous, with the angle rounded. The skin is covered with small rough scales, each of which bears a well developed spinate projection, which consists of a central spine from which radiate four compressed wings, each one terminating at its outer angle in a somewhat shorter spine than the central one.

Color.—Uniform sandy brown.

The Shark described above was sent to the Australian Museum, by the authorities of the Tasmanian Museum, Hobart, for identification and preservation, and was placed in my hands for description previous to being returned. The specimen was picked up on the shore of Bruny Island, Tasmania, in a dried state, but on being relaxed was found to be in a fair state of preservation. The enormous height of the dorsal fins, and their contiguity, the one to the other, separates this species at a glance from *C. salviani*; the scales also differ considerably.

Type.—In the Tasmanian Museum.

The specific name has been given to it at the request of Mr. Alex. Morton, Curator of the Tasmanian Museum.