

AUSTRALIAN MUSEUM SCIENTIFIC PUBLICATIONS

North, A. J., 1899. Ornithological notes. IX. Description of the nest and eggs of *Micraeca pallida*, De Vis. *Records of the Australian Museum* 3(5): 107. [17 April 1899].

doi:10.3853/j.0067-1975.3.1899.1149

ISSN 0067-1975

Published by the Australian Museum, Sydney

nature culture **discover**

Australian Museum science is freely accessible online at
<http://publications.australianmuseum.net.au>
6 College Street, Sydney NSW 2010, Australia



For the sake of comparison the following measurements are given

	Total length of skin.	Wing.	Tail.	Bill.	Tarsus.	Localities.
<i>P. leilavalensis</i> , ad. sk	6 inches	2·8	2·7	0·42	0·75	Fullerton River, N. Queensland.
<i>P. penicillata</i> ♂ ad. sk.	6·7 „	3·45	3·3	0·5	0·82	Near Adelaide, S. Australia.
„ ♂ ad. sk.	6·6 „	3·3	3·2	0·47	0·8	Bourke, Western N. S. Wales.
„ ♂ ad. sk.	6·5 „	3·3	3·1	0·47	0·8	Dawson River, Queensland.
<i>P. flavescens</i> , ♂ ad. sk.	5·7 „	2·95	2·7	0·5	0·7	Georgetown, Gulf District.
„ ♀ ad. sk.	5·4 „	2·92	2·7	0·48	0·7	Derby, N. West Australia.

IX.—DESCRIPTION OF THE NEST AND EGGS OF *MICRÆCA* *PALLIDA*, DE VIS.

Dr. W. Macgillivray has kindly forwarded the following description of the nest of this species, together with the eggs and skin of the bird for identification.

“Two nests of this species of *Micræca* were taken by my brother Mr. A. S. Macgillivray on Leilavale Station, Fullerton River, North Queensland, between the 20th and 25th December, 1897. They were built rather low down on horizontal branches in a patch of *Giddea* scrub and each contained a pair of eggs. A nest my brother sent was slightly smaller but more substantially built than that of *M. fascinans*, and of much the same material, the outside being ornamented with bits of bark and lichen attached by means of cobweb.”

The eggs are oval in form, one specimen having a pale bluish-grey ground colour, which is freckled and spotted with faint purple and purplish-brown, predominating and becoming darker as usual on the thicker end of the shell; the other is of a warm stone white ground colour, and in many places the markings which are of a light reddish-purple are confluent and intermingled with faint underlying spots of pale greyish-lilac. Length (A) 0·69 x 0·55 inch; (B) 0·67 x 0·56 inch.

The range of this species probably extends right across the northern portion of the Australian Continent, for there are specimens in the Museum, obtained by Mr. E. J. Cairn at Derby, North-west Australia in 1886.