

# AUSTRALIAN MUSEUM SCIENTIFIC PUBLICATIONS

Whitelegge, T., 1897. On *Stichopus mollis*, Hutton. *Records of the Australian Museum* 3(2): 50. [5 August 1897].

doi:10.3853/j.0067-1975.3.1897.1129

ISSN 0067-1975

Published by the Australian Museum, Sydney

nature culture **discover**

Australian Museum science is freely accessible online at  
<http://publications.australianmuseum.net.au>  
6 College Street, Sydney NSW 2010, Australia



Gould's *C. parvissima* is a decidedly smaller race, and is further distinguished by its upper parts being more strongly washed with olive. A specimen from Cape York measures, total length 6·3 inches, wing 3·5, tail 2·7, bill 0·75. It ranges as far south as the Herbert River. The wing-measurement varies from 3·5 to 3·6 inches. In a large series of specimens examined I can find no gradation in size between *C. rufigaster* and its smaller northern ally, *C. parvissima*. Dr. Sharpe, in the "Catalogue of Birds in the British Museum,"\* has described a specimen of the latter race from Cape York under the name of the larger species, *Pinarolestes rufigaster*, Gould.

### ON *STICHOPUS MOLLIS*, HUTTON.

By THOMAS WHITELEGGE, Zoologist, Australian Museum.

DURING a recent visit to Eden, Dr. J. C. Cox obtained a *Holothurian*, which he presented to the Museum collection.

The species proves to be *Stichopus mollis*, Hutton, hitherto only recorded from New Zealand. At first I referred it to *S. sordidus*, Th  el,† but on consulting a paper by Prof. A. Dendy‡ since received, I find that the "Challenger" specimens are considered to be examples of Hutton's *Holothuria mollis*. I agree with the opinion as expressed by Prof. Dendy. There appears to be no character to distinguish them except that of colour, which is evidently variable.

The colour of the Eden example in formol is light yellowish-brown, with the disks of the pedicels and the tips of the dorsal papill  e darker.

The large bilateral plates form a ring on the margin of the disk and encircle the large central perforate plate at the extremity of the pedicel.

The dorsal papill  e are supported by curved, smooth, spiny, or branched rods, disposed transversely; the lower ones tend to form plates similar to those of the ventral pedicels; apically each papilla terminates in a large perforate plate, which is surrounded by a series of stout moniliform rods, with either simple or spinose ends.

Length of specimen 130 mm.

\* Brit. Mus. Cat. Birds iii. 1877, p. 296.

† Chall. Rep. Zool., xiv., p. 162, pl. viii., fig. 3.

‡ Jour. Linn. Soc. Zool. xxvi., 1897, p. 46, pl. vii., figs. 73-82.

5th August, 1897.