## SOME NATICOIDS FROM QUEENSLAND.

#### By

### CHARLES HEDLEY.

## (Plate xxii.)

The identity of a series of tropical species of Naticoids is unsatisfactory because of defective treatment in Reeve's monograph of *Natica* in the "Conchologica Iconica," which has governed subsequent studies of this group.

To put taxonomy on a better basis, I have here reviewed the nomenclature of several obscure species, and to facilitate their recognition also present figures of them.

The first name given to a group, resembling the Linnean Nerita mammilla, of polished massive white or yellow shells, was Humphrey's Uber. Uber flemingianum Recluz, jukesii Reeve, labyrintheum Hedley, mellosum Hedley and pyriforme Recluz, can be associated together.

Those species associated with *U. pes-elephantis*, being large, massive shells with a wide umbilicus containing a stout spiral funicle, may be grouped under *Mammillaria*, Swainson; namely, *U. citrinum* Philippi, columnare Recluz, powisianum Recluz, and pes-elephantis Deshayes.

Lastly, those thinner ovate and variegated shells grouped round *Uber mammatum* can be ranked under *Mammilla* Schumacher; such are *Uber mammatum* Bolten, *melanostomoides* Quoy & Gaimard, *nuxcastaneum* Martyn, *opacum* Recluz, *sebæ* Recluz and *simiæ* Deshayes.

# UBER Humphrey.

- Uber Humphrey, Mus. Calonnianum, 1797, p. 21, first species and type Nerita mammilla Linné.
- Albula Bolten Mus. Bolt. (2), 1798, p. 20, first species Albula albumen Linné (not Albula Gronovius, Zooph., i, 1763, p. 102).
- Mammilla Schumacher Essai, 1817, p. 190, type Mammilla fasciata Schumacher.
- Polinices Montfort Conch. Syst. ii, 1810, p. 222, type Polinices albus Montfort.
- Mammillaria Swainson, Treat. Malac., 1840, p. 345, type M. lactea Swainson.
- Naticaria Swainson, Treat. Malac, 1840, p. 345, type N. melanostoma Mart.